


Myrbetriq (Mirabergron) in Patients with Overactive Bladder (OAB)

Purpose: Evaluate appropriateness of Myrbetriq utilization in patients with overactive bladder

Treatment Plan for Overactive Bladder



Drug for Overactive Bladder

Average Cost Per Day

Myrbetriq (Mirabergron) ER	\$15.36
Myrbetriq (Mirabergron)	\$13.35
Gemtesa	\$13.87
Trospium chloride ER	\$3.65
Trospium chloride	\$0.63
Tolterodine tartrate	\$1.27
Solifenacin succinate	\$0.17
Oxybutynin chloride ER	\$0.15
Oxybutynin chloride	\$0.15

Continued on back →

Myrbetriq vs Oxybutynin

Myrbetriq (Mirabegron)

Used for overactive bladder with symptoms of urge urinary incontinence, urgency and urinary frequency

Onset of action is ~4 to 6 weeks

Main side effects are hypertension, nasopharyngitis, urinary tract infection and headaches

Much higher cost

No generic available

Oxybutynin (Generic)

Used for relief of symptoms of overactive bladder or symptoms of detrusor overactivity

Onset of action ~4 weeks

Main side effects of dry mouth, nausea, blurred vision, and headache (anticholinergic effects)

Cost savings option

Generic

If you have any additional questions, please contact Claire Barlow, Clinical Pharmacist, at claire.barlow@commonspirit.org.

Disclaimer: This information is not intended to dictate or substitute your professional judgment in a particular patient.

For more info,
visit our website

